

1D12

Stark and Zeeman deceleration of atoms and molecules

S. D. Hogan, A. W. Wiederkehr, H. Schmutz, M. Andrist and F. Merkt
Laboratorium für Physikalische Chemie, ETH Zurich, Switzerland

In the past years considerable efforts have been invested to develop general methods with which to produce (ultra-)cold samples of atoms and molecules that cannot be laser cooled. Methods developed so far include the photoassociation of ultracold atoms [1,2], buffer-gas cooling [3] and multistage Stark deceleration [4]. With the aim of performing very high resolution spectroscopy we have recently contributed to the development of two new methods of producing cold samples of atoms and molecules starting from supersonic beams: Rydberg-Stark deceleration [5,6] and multistage Zeeman deceleration [7,8]. The talk will provide a detailed description of these two methods.

The former method exploits the very large dipole moments (more than 100 Debye) that can be induced in atomic and molecular Rydberg states by electric fields and was recently used to stop and trap electrostatically clouds of translationally cold Rydberg atoms after deceleration in a single-stage device [9]. The latter method exploits the Zeeman effect in paramagnetic species and the ability to switch on and off large magnetic fields (up to 2 Tesla) in less than 5 μ s. It was used recently to decelerate an atomic sample initially in a supersonic beam to zero velocity in the laboratory reference frame and subsequently load the atoms into a magnetic trap. The deceleration methods, the diagnostic methods to characterize the velocity distributions of the decelerated species and the trapping methods will be illustrated by experiments conducted on atomic argon and hydrogen.

[1] A. Fioretti, D. Comparat, A. Crubellier, O. Dulieu, F. Masnou-Seeuws and P. Pillet, *Phys. Rev. Lett.* **80**, 4402 (1998)

[2] M. Yasuda, T. Kishimoto, M. Takamoto and H. Katori, *Phys. Rev. A* **73**, 011403(R) (2006)

[3] J. M. Doyle, B. Friedrich, J. Kim and D. Patterson, *Phys. Rev. A* **52**, 2515 (1995)

[4] H. L. Bethlem, G. Berden and G. Meijer, *Phys. Rev. Lett.* **83**, 1558 (1999).

[5] S. R. Procter, Y. Yamakita, F. Merkt and T. P. Softley, *Chem. Phys. Lett.* **374**, 667 (2003)

[6] E. Vliegen, H. J. Wörner, T. P. Softley and F. Merkt, *Phys. Rev. Lett.* **92**, 033005 (2004)

[7] N. Vanhaecke, U. Meier, M. Andrist, B. H. Meier and F. Merkt, *Phys. Rev. A* **75**, 031402(R) (2007)

[8] S. D. Hogan, A. W. Wiederkehr, M. Andrist, H. Schmutz and F. Merkt, *J. Phys. B* **41**, 081005 (2008)

[9] S. D. Hogan and F. Merkt, *Phys. Rev. Lett.* **100**, 043001 (2008)